

ON CURVATURES OF HOMOGENEOUS FINSLER-KROPINA SPACE

KERIYAPPA CHANDRU¹ AND SENAJJI KAMPLAPPA NARASIMHAMURTHY^{2*}

ABSTRACT. In Finsler geometry, one of the central problem is to study curvature properties of special class of Finsler spaces such as homogeneous-Finsler spaces. In this paper, we give an explicit formula of the S-curvature of homogeneous Finsler space with Kropina metric and proved that a homogeneous Kropina space with almost isotropic S-curvature must have vanishing S-curvature. We also derived an explicit formula of the mean Berwald curvature E_{ij} of homogeneous Finsler-Kropina space.

1. INTRODUCTION

Finsler geometry is just the Riemannian geometry without quadratic restriction [3]. Thus it can be considered as the nonlinear version of Riemannian geometry, but nonlinear characterization of the metrics makes Finsler geometry much more complicated and difficult compared to Riemannian geometry [2]. However, recently this geometry has been studied extensively and has proven to be very use full in Physics and Biology [3]. One of the central problem in Finsler geometry is to study curvature properties of special class of Finsler spaces such as homogeneous-Finsler spaces.

Let (M, F) be a Finsler space, where M be the connected smooth manifold and F be the Finsler metric. (α, β) -metrics are the class of Finsler metrics, for example Randers metric, Kropina metric, Matsumoto metric, etc. Kropina metric is of the form $F = \frac{\alpha^2}{\beta}$, where α is Riemannian metric and β is the one form. Kropina metrics were first introduced by L. Berwald in connection with a two-dimensional Finsler space with rectilinear extremal and were investigated by V. K. Kropina [14].

A Finsler-Kropina space (M, F) becomes homogeneous if the group of isometries $I(M, F)$ of (M, F) acts transitively on M . As described in [7], for homogeneous Finsler space, M can be written as a coset space G/H with (α, β) -metric of the form $F = \alpha\phi(s)$, where $s = \frac{\beta}{\alpha}$ with α a G -invariant Riemannian metric on G/H and β a G -invariant vector field on G/H . Therefore the Lie algebra $\mathfrak{g}$ of G

Date: Received: Apr 27, 2016; Accepted: Aug 10, 2016.

* Corresponding author.

2010 *Mathematics Subject Classification.* Primary 53B40; Secondary 53C60.

Key words and phrases. Finsler space, (α, β) -metric, Homogeneous Finsler space, S-curvature, Mean-Berwald curvature.

expressed as composition of h and m . i.e.,

$$g = h + m \text{ (direct sum of subspaces)} \quad (1.1)$$

such that $Ad(h)(m) \subset m$, $h \in H$ and we can identify M with the tangent space of (G/H) at the origin $o = H$. Further, the one form β corresponds to a vector field $\bar{X}$ on G/H which is generated by $Ad(H)$ -invariant vector in m with length < 1 .

The goal of this paper is to derive an explicit formula for the S -curvature of homogeneous Finsler space with Kropina metric. The notion of S -curvature for a Finsler space introduced by Z. Shen in [15]. It is a quantity to measure the rate of change of the volume form of a Finsler space along the geodesics. Recently many geometers studied curvature properties of homogeneous Finsler space [7, 8]. In 2007, S. Deng and Z. Hou studied that a homogeneous Finsler spaces with non-positive flag curvature [8]. In 2009, S. Deng obtained the explicit formula of S -curvature of homogeneous Randers space with almost isotropic S -curvature must have vanishing S -curvature [9]. As for reference, there is an explicit formula for S -curvature in a local standard coordinate system by Z. Shen [15]. However, for a homogeneous spaces there should have a formula which does not use local coordinates. In this article we are studied curvature properties of homogeneous Finsler-Kropina space and find the formula for S -curvature and mean-Berwald curvature.

In the following, we shall use Einstein summation convention unless otherwise stated.

2. PRELIMINARIES

In this section we recall some known results and notations. Finsler space is a smooth manifold possessing a Finsler metric. A formal definition of a Finsler space is as follows:

Definition 2.1. A Finsler space is a triple $F^n = (M, D, F)$, where M is an n -dimensional manifold, D is an open subset of a tangent bundle TM and F is a Finsler metric defined as a function $F : TM \rightarrow [0, \infty)$, with the following properties:

- (1) Regular: F is C^∞ on the entire tangent bundle $TM \setminus \{0\}$.
- (2) Positive homogeneous: $F(x, \lambda y) = \lambda F(x, y)$, $\forall \lambda \geq 0$.
- (3) Strong convexity: The $n \times n$ Hessian matrix

$$g_{ij} = \frac{1}{2} [F^2]_{y^i y^j}$$

is positive definite at every point on $TM \setminus \{0\}$, where $TM \setminus \{0\}$ denotes the tangent vector y is non zero in the tangent bundle TM .

In Finsler geometry, (α, β) -metrics is a important class of Finsler metrics [10]. A Finsler metric $F = F(x, y)$ is said to be (α, β) -metric if it can be expressed in

the form,

$$F = \alpha\phi(s), \quad s = \frac{\beta}{\alpha}, \quad (2.1)$$

where $\alpha = \sqrt{g_{ij}(x)y^i y^j}$ is a Riemannian metric and $\beta = b_i(x)y^i$ is a 1-form. Here $\phi = \phi(s)$ is a C^∞ positive function on an open interval $(-b_0, b_0)$ satisfying

$$\phi(s) - s\phi'(s) + (b^2 - s^2)\phi''(s) > 0, \quad |s| \leq 0 < b_0, \quad b = \|\beta(x)\|_\alpha. \quad (2.2)$$

In this article we considered an important example of such (α, β) -metrics is Kropina metric introduced by V. K. Kropina [12], expressed in the following form,

$$\begin{aligned} F &= \frac{\alpha(x, y)^2}{\beta(x, y)} \\ &= \alpha\phi(s); \quad \phi(s) = \frac{1}{s}, \end{aligned}$$

where $s = \frac{\beta}{\alpha}$, $\alpha = \sqrt{g_{ij}(x)y^i y^j}$ is the Riemannian metric and $\beta = b_i(x)y^i$ is the 1-form.

Let

$$r_{ij} = \frac{1}{2}(b_{i|j} + b_{j|i}), \quad s_{ij} = \frac{1}{2}(b_{i|j} - b_{j|i}). \quad (2.3)$$

Clearly, β is closed if and only if $s_{ij} = 0$. We put

$$\begin{aligned} r_{i0} &= r_{ij}y^j, \quad r_{00} = r_{ij}y^i y^j, \quad r_j = b^i r_{ij}, \\ s_{i0} &= s_{ij}y^j, \quad s_j = b^i s_{ij}, \quad r_0 = r_j y^j, \quad s_0 = s_j y^j. \end{aligned}$$

For a Finsler metric $F = F(x, y)$, the geodesics are characterized by system of second order ordinary differential equations,

$$\ddot{x}^i + 2G^i(x, \dot{x}) = 0,$$

where $G^i = G^i(x, y)$ are called spray coefficients and are given by,

$$G^i = y^i \frac{\partial}{\partial x^i} - 2G^i(x, y) \frac{\partial}{\partial x^i} \quad (2.4)$$

in standard local coordinates (x^i, y^i) in TM , then the spray of F is the vector field $G = y^i \frac{\partial}{\partial x^i} - 2G^i(x, y) \frac{\partial}{\partial y^i}$ [10].

The S -curvature is an important non-Riemannian quantity introduced by Z. Shen [15], defined with respect to a volume form $dV = \sigma(x)dx$ by,

$$S = \frac{\partial G^m}{\partial y^m} - y^m \frac{\partial}{\partial x^m} (\ln \sigma_F(x)). \quad (2.5)$$

Here the volume form can be the Busemann-Hausdorff volume form $dV_{BH} = \sigma_{BH}(x)dx$, where

$$\sigma_{BH}(x) = \frac{Vol(B^n)}{Vol\{(y^i) \in R^n \mid F(x, y) < 1\}}$$

or the Holmes-Thompson volume form $dV_{HT} = \sigma_{HT}(x)dx$, where

$$\sigma_{HT}(x) = \frac{1}{Vol(B^n)} \int_{F(x, y) < 1} \det(g_{ij}(x, y)) dy.$$

Unless specified, the S -curvature usually defined with respect to the Busemann-Hausdorff volume form.

Definition 2.2. A Finsler space (M, F) is said to have almost isotropic S -curvature if there exists a smooth function $c(x)$ on M and a closed 1-form η such that:

$$S(x, y) = (n + 1)c(x)F(y) + \eta(y), \quad x \in M, \quad y \in T_x M. \quad (2.6)$$

Note:(1). F is of isotropic S -curvature if $c = c(x)$ is a scalar for and $\eta = 0$.
 (2). F is of constant S -curvature if c is a constant and $\eta = 0$.

In [12], authors defined the S -curvature of the (α, β) -metric $F = \alpha\phi(s)$, in local coordinate system by,

$$S = \left\{ 2\Psi - \frac{f'(b)}{bf(b)} \right\} (r_0 + s_0) - \alpha^{-1} \frac{\Phi}{2\Delta^2} (r_{00} - 2\alpha Q s_0), \quad (2.7)$$

where

$$Q = \frac{2 + 2s}{1 - s^2}, \quad \Delta = 1 + sQ + (b^2 - s^2)Q', \quad \Psi = \frac{Q'}{2\Delta} \quad (2.8)$$

$$\Phi = -(Q - sQ') \{n\Delta + 1 + sQ\} - (b^2 - s^2)(1 + sQ)Q'' \quad (2.9)$$

and the function $f(b)$ in the formula is defined as follows,

$$f(b) = \begin{cases} \frac{\int_0^\pi \sin^{n-2} t dt}{\int_0^\pi \frac{\sin^{n-2} t}{\phi(b \cos t)^n} dt} & \text{if } dV = dV_{BH}; \\ \frac{\int_0^\pi (\sin^{n-2} t) T(b \cos t) dt}{\int_0^\pi \sin^{n-2} t dt} & \text{if } dV = dV_{HT}, \end{cases}$$

where $T(s) = \phi(\phi - s\phi')^{n-2} \{(\phi - s\phi') + (b^2 - s^2)\phi''\}$. Then the volume form dV is given by $dV = f(b)dV_\alpha$, where $dV_\alpha = \sqrt{\det(\alpha_{ij})}dx$, denote the Riemannian volume form of α .

3. S -CURVATURE OF HOMOGENEOUS FINSLER-KROPINA SPACE

Recall that two Finsler spaces (M_1, F_1) and (M_2, F_2) are said to be isometric if there exists a diffeomorphism φ from M_1 onto M_2 , such that $F_1(x, y) = F_2(\varphi(x), \varphi(y))$ for any $x_1 \in M_1$ and $y \in T_x M_1$. In 2006 [7], S. Deng and Z. Hou proved that the group of isometries of Finsler space is a Lie transformation group on the original manifold which can be used to study homogeneous Finsler spaces.

A homogeneous Finsler-Kropina space M can be written as a coset space G/H with a G -invariant Kropina metric $F = \frac{\alpha^2}{\beta}$, where both Riemannian metric α and the one form β are invariant under the action of G . In [9], S. Deng and Z. Hou, specified that, β corresponding to a unique vector u in $T_o(G/H)$ which is fixed under linear isotropy representation of H on $T_o(G/H)$ and $o = H$ is the origin of G/H . And in [8], S. Deng and X. Wang gives the following result,

Theorem 3.1. [8] *Let $F = \alpha\phi(s)$ be a G -invariant (α, β) metric on the reduction homogeneous manifold G/H with a decomposition of the Lie algebra*

$$g = h + m. \quad (3.1)$$

Then the S -curvature of F has the form

$$S(o, y) = -\frac{1}{\alpha(y)} \frac{\Phi}{2\Delta^2} \{-c\langle [u, y]_m, y \rangle - \alpha(y)Q\langle [u, y]_m, u \rangle\}, \quad y \in m, \quad (3.2)$$

where u is the vector in m corresponding to the 1-form β , and we have identified m with the tangent space of G/H at origin $o = H$.

In this section according to the above theorem (3.1) we find the formula of S -curvature for homogeneous Finsler space with Kropina metric. By definition (2.1), we can write Kropina metric $F = \frac{\alpha^2}{\beta}$ as $F = \alpha\phi(s)$, where $\phi(s) = \frac{1}{s}$. Now, for this Kropina metric by the equations (2.8) and (2.9) we have,

$$Q = -\frac{1}{2s}, \quad Q' = \frac{1}{2s^2}, \quad (3.3)$$

$$\Delta = \frac{b^2}{2s^2}, \quad \Phi = \frac{(n+1)b^2}{2s^3}. \quad (3.4)$$

Since $(G/H, F)$ is homogeneous, it is enough to compute the S -curvature at the origin $o = H$. Let $(U, (x^1, x^2, \dots, x^n))$ be the local coordinate system. With reference to [4], we have to find the S -curvature in local coordinate system, for that we need to evaluate the following quantities at the origin:

- (1) $r_{ij} = \frac{1}{2}(b_{i;j} + b_{j;i})$ and b_i 's are defined by $\beta = b_i(x)dx^i$.
- (2) $s_i = b_j s_i^j$ and s_j^i are defined by $s_j^i = \frac{1}{2} \left(\frac{\partial b_i}{\partial x^j} - \frac{\partial b_j}{\partial x^i} \right)$.
- (3) $s_0 = s_i y^i$
- (4) $\rho_0 = \rho_{xi} y^i$, where $\rho = \sqrt{1 - \|\beta\|}$ and $\|\beta\|$ is the length of the 1-form β with respect to α .

Let $\langle, \rangle$ be the corresponding inner product on m . Then we have to consider the Levi-Civita connection of $(G/H, \alpha)$ which will be useful to calculate the S -curvature of homogeneous Finsler-Kropina spaces. There are many versions of the formula of the connection for Killing vector fields. But, we are working on the differential of (left) invariant vector fields on G/H and we adopt the formula in [8]. Now for $v \in g$, define a one parameter transformation group φ_t , $t \in R$ of G/H by

$$\varphi_t(gH) = (\exp(tv)g)H, \quad g \in G.$$

Then φ_t generates a vector field on G/H which is a Killing vector field (this is called the fundamental vector field generated by v in [8]). We denote this vector field by $\bar{v}$. In [14] authors defined the following formula,

$$\langle \nabla_{v_1} v_2 |_o, w \rangle = \frac{1}{2} (-\langle [v_1, v_2]_m, w \rangle + \langle [w, v_2]_m, v_1 \rangle + \langle [w, v_1]_m, v_2 \rangle), \quad v_1, v_2, w \in m. \quad (3.5)$$

where $o = H$ is the origin of the coset space and $[v_1, v_2]_m$ denote the projection of $[v_1, v_2]$ to m corresponding to the decomposition (3.1).

Now by using the formula(2.3), we have to compute r_{00} and s_0 . First consider,

$$b_i = \beta \left(\frac{\partial}{\partial x^i} \right) = \langle \tilde{u}, \frac{\partial}{\partial x^i} \rangle = c \langle \frac{\partial}{\partial x^n}, \frac{\partial}{\partial x^i} \rangle,$$

Then by differentiating b_i we get,

$$\begin{aligned} \frac{\partial b_i}{\partial x^j} &= c \frac{\partial}{\partial x^j} \langle \frac{\partial}{\partial x^n}, \frac{\partial}{\partial x^i} \rangle \\ &= c \left(\langle \nabla_{\frac{\partial}{\partial x^j}} \frac{\partial}{\partial x^n}, \frac{\partial}{\partial x^i} \rangle + \langle \frac{\partial}{\partial x^n}, \nabla_{\frac{\partial}{\partial x^j}} \frac{\partial}{\partial x^i} \rangle \right). \end{aligned} \quad (3.6)$$

Hence at the origin we have (here we use the symmetry of the connection: $\nabla_{\frac{\partial}{\partial x^j}} \frac{\partial}{\partial x^i} - \nabla_{\frac{\partial}{\partial x^i}} \frac{\partial}{\partial x^j} - [\frac{\partial}{\partial x^i}, \frac{\partial}{\partial x^j}] = 0$),

$$s_{ij}(o) = \frac{1}{2} c \left(\langle \nabla_{\frac{\partial}{\partial x^n}} \frac{\partial}{\partial x^j}, \frac{\partial}{\partial x^i} \rangle - \langle \nabla_{\frac{\partial}{\partial x^n}} \frac{\partial}{\partial x^i}, \frac{\partial}{\partial x^j} \rangle \right) |_o \quad (3.7)$$

Then by equation (3.5) we have,

$$s_{ij}(o) = \frac{1}{2} \langle [u_i, u_j]_m, u_n \rangle. \quad (3.8)$$

Since at the origin we have $(a_{ij}) = I_n$, we get

$$s_i^i(o) = a^{ik}(o) s_{kj}(o) = \sum_{k=1}^n \delta_i^k s_{kj}(o) = s_{ij}(o).$$

Therefore

$$s_i(o) = b_l(o) s_i^l(o) = c s_i^n(o) = s_{ni}(o). \quad (3.9)$$

Now for $y = y^i u_i \in m$, we have

$$\begin{aligned} s_0(y) &= y^l s_l(o) = c y^l s_{nl}(o) = \frac{1}{2} y^l \langle [u_n, u_l]_m, u_n \rangle \\ &= \frac{1}{2} \langle [c u_n, y^l u_l]_m, c u_n \rangle \\ &= \frac{1}{2} \langle [u, y]_m, u \rangle. \end{aligned} \quad (3.10)$$

Next we compute r_{ij} . Suppose $i \geq j$. Then we have

$$\begin{aligned} r_{ij}(o) &= \frac{1}{2} (b_{i;j} + b_{j;i}) |_o \\ &= \frac{1}{2} \left(\frac{\partial b_i}{\partial x^j} - b_l \Gamma_{ji}^l + \frac{\partial b_j}{\partial x^i} - b_l \Gamma_{ji}^l \right) |_o \\ &= \frac{1}{2} \left(\frac{\partial b_i}{\partial x^j} + \frac{\partial b_j}{\partial x^i} \right) |_o - c \Gamma_{ji}^l(o). \end{aligned} \quad (3.11)$$

By (3.6) we have

$$\frac{1}{2} \left(\frac{\partial b_i}{\partial x^j} + \frac{\partial b_j}{\partial x^i} \right) |_o = -\frac{1}{2} \langle [u_i, u_j]_m, u_n \rangle, \quad i \geq j. \quad (3.12)$$

Combining the above equation (3.12) with the connection coefficients Γ_{ij}^l defined in [9], we get

$$r_{ij}(o) = -\frac{1}{2}c(\langle [u_n, u_j]_m, u_i \rangle + \langle [u_n, u_i]_m, u_j \rangle), \quad i \geq j. \quad (3.13)$$

Further

$$r_{00}|_o = r_{ij}(o)y^i y^j = -c\langle [u_n, y]_m, y \rangle. \quad (3.14)$$

By substituting the above quantities and the values of (3.3) and (3.4) into the equation (3.2) we get the formula for S -curvature of homogeneous Finsler-Kropina metric at the origin as follows,

$$S(o, y) = -\frac{(n+1)s}{b^2\alpha(y)} \left\{ -c\langle [u, y]_m, y \rangle + \frac{\alpha(y)}{2s} \langle [u, y]_m, u \rangle \right\}. \quad (3.15)$$

Summarizing above statements we get,

Theorem 3.2. *Let $F = \alpha\phi(s)$ be a G -invariant (α, β) -metric on the reductive homogeneous manifold G/H , with a decomposition of the Lie algebra*

$$g = h + m. \quad (3.16)$$

Then the S -curvature of $F = \frac{\alpha^2}{\beta}$ has the form

$$S(o, y) = -\frac{(n+1)s}{b^2\alpha(y)} \left\{ -c\langle [u, y]_m, y \rangle + \frac{\alpha(y)}{2s} \langle [u, y]_m, u \rangle \right\}, \quad (3.17)$$

where u is the vector in m corresponding to the 1-form β , and we have identified m with the tangent space of G/H at the origin $o = H$.

As a direct application of the above formula we get,

Theorem 3.3. *Let $(G/H, F)$ be as in the above theorem (3.2). Then homogeneous Kropina metric F has isotropic S -curvature if and only if F has vanishing S -curvature.*

Proof: By using the formula derived in the previous theorem (3.2), it is enough to show the direct part. So, suppose F has isotropic S -curvature;

$$S(x, y) = (n+1)c(x)F(y), \quad x \in G/H, \quad y \in T_x(G/H). \quad (3.18)$$

Letting $x = 0$ and $y = u$ then by (3.17) we get $c(0) = 0$. Hence $S(o, y) = 0, \forall y \in T_o(G/H)$. Since F is a homogeneous metric, we must have $S = 0$ everywhere.

4. MEAN BERWALD CURVATURE OF HOMOGENEOUS FINSLER-KROPINA SPACE

One more important non-Riemannian quantity in Finsler geometry is mean-Berwald curvature. In this section we apply the results of Section 3 to give a formula of mean Berwald curvature of homogeneous Finsler-Kropina metrics. In [6], X. Cheng and Z. Shen, give the mean Berwald curvature for Finsler metrics as follows,

$$E_{ij} = \frac{1}{2} \frac{\partial^2}{\partial y^i \partial y^j} \left(\frac{\partial G^m}{\partial y^m} \right), \quad (4.1)$$

where $G^m = G^m(x, y)$ are the spray coefficients. By (2.5), S -curvature can be defined as

$$S = \frac{\partial G^m}{\partial y^m} - (\ln \sigma(x))_{x^k} y^k. \quad (4.2)$$

Here, since $\sigma(x)$ is the function of x , the second term in the above equation $(\ln \sigma(x))$ is also function of x only. Therefore,

$$\frac{\partial^2}{\partial y^i \partial y^j} [(\ln \sigma(x))_{x^k} y^k] = 0.$$

Hence, the formula for mean Berwald curvature obtained by direct computation,

$$\begin{aligned} \frac{\partial^2 S}{\partial y^i \partial y^j} &= \frac{\partial^2}{\partial y^i \partial y^j} \left[\frac{\partial G^m}{\partial y^m} - (\ln \sigma(x))_{x^k} y^k \right] \\ &= \frac{\partial^2}{\partial y^i \partial y^j} = 2E_{ij}. \end{aligned} \quad (4.3)$$

So that, clearly by partially differentiating the expression of S -curvature derived in section 3, we can determine the formula for mean Berwald curvature of the homogeneous Finsler-Kropina space. i.e.,

$$2E_{ij} = \frac{\partial^2 S}{\partial y^i \partial y^j}. \quad (4.4)$$

Before the computation, we consider the identities,

$$\frac{\partial^2 s}{\partial y^m} = \frac{1}{\alpha} \left(b_m - s \frac{y_m}{\alpha} \right), \quad \frac{\partial \alpha}{\partial y^m} = \frac{y^m}{\alpha}. \quad (4.5)$$

Theorem 4.1. *Let $(G/H, F)$ be a homogeneous Finsler-Kropina space with the metric of the form $F = \frac{\alpha^2}{\beta}$ where α be a G -invariant Riemannian metric on G/H and β be a G -invariant vector field on G/H . Then the mean Berwald curvature of homogeneous Finsler-Kropina space is of the form*

$$E_{ij} = \frac{(n+1)}{2b^2} \left\{ \frac{s}{\alpha(y)} [\langle [u, u_i]_m, u_j \rangle + \langle [u, u_j]_m, u_i \rangle] + \langle [u, y]_m, y \rangle A \right\}, \quad (4.6)$$

where A is defined in equation (4.17).

Proof: Let us consider Finsler-Kropina metric $F = \alpha\phi(s)$, where $\phi(s) = \frac{1}{s^2}$. Put $\psi = \phi - s\phi$, then we have,

$$Q = \frac{\phi'}{\psi} = -\frac{1}{2s}, \quad Q' = \frac{1}{2s^2}, \quad \Delta = 1 + sQ + (b^2 - s^2)Q' = \frac{b^2}{2s^2}, \quad (4.7)$$

$$\Phi = -(Q - sQ')n\Delta + 1 + sQ - (b^2 - s^2)(1 + sQ)Q'' = \frac{(n+1)b^2}{2s^3}. \quad (4.8)$$

Letting $c = 1$ in the formula of S -curvature defined in equation (3.17), we get

$$\begin{aligned} S(o, y) &= -\frac{(n+1)s}{b^2\alpha(y)} \left\{ -\langle [u, y]_m, y \rangle + \frac{\alpha(y)}{2s} \langle [u, y]_m, u \rangle \right\}, \\ &= \frac{(n+1)}{b^2} \left\{ \frac{s}{\alpha(y)} \langle [u, y]_m, y \rangle - \frac{1}{2} \langle [u, y]_m, u \rangle \right\}. \end{aligned} \quad (4.9)$$

Now by using (4.4) we obtain,

$$\begin{aligned}
2E_{ij} &= \frac{\partial^2 S(o, y)}{\partial y^i \partial y^j}, \\
&= \frac{(n+1)}{b^2} \frac{\partial^2}{\partial y^i \partial y^j} \left\{ \frac{s}{\alpha(y)} \langle [u, y]_m, y \rangle - \frac{1}{2} \langle [u, y]_m, u \rangle \right\}, \\
&= \frac{(n+1)}{b^2} \left\{ \frac{\partial^2}{\partial y^i \partial y^j} \left(\frac{s}{\alpha(y)} \langle [u, y]_m, y \rangle \right) - \frac{\partial^2}{\partial y^i \partial y^j} \left(\frac{\langle [u, y]_m, u \rangle}{2} \right) \right\} \quad (4.10)
\end{aligned}$$

Now consider the first term,

$$\begin{aligned}
\frac{\partial^2}{\partial y^i \partial y^j} \left\{ \frac{s}{\alpha(y)} \langle [u, y]_m, y \rangle \right\} &= \frac{s}{\alpha(y)} \frac{\partial^2}{\partial y^i \partial y^j} (\langle [u, y]_m, y \rangle) \\
&\quad + \langle [u, y]_m, y \rangle \frac{\partial^2}{\partial y^i \partial y^j} \left(\frac{s}{\alpha(y)} \right), \quad (4.11)
\end{aligned}$$

but we get,

$$\begin{aligned}
\frac{\partial \langle [u, y]_m, y \rangle}{\partial y^i} &= \langle [u, u_i]_m, y \rangle + \langle [u, y]_m, u_i \rangle, \\
\frac{\partial^2 \langle [u, y]_m, y \rangle}{\partial y^i \partial y^j} &= \langle [u, u_i]_m, u_j \rangle + \langle [u, u_j]_m, u_i \rangle, \quad (4.12)
\end{aligned}$$

and

$$\frac{\partial^2}{\partial y^i \partial y^j} \left(\frac{s}{\alpha(y)} \right) = \frac{8s y_i y_j}{\alpha^5} - \frac{2}{\alpha^4} (b_j y_i + b_i y_j) - \frac{2s}{\alpha^3} \delta_i^j. \quad (4.13)$$

Therefore we have,

$$\begin{aligned}
\frac{\partial^2}{\partial y^i \partial y^j} \left\{ \frac{s}{\alpha(y)} \langle [u, y]_m, y \rangle \right\} &= \frac{s}{\alpha(y)} [\langle [u, u_i]_m, u_j \rangle + \langle [u, u_j]_m, u_i \rangle] \\
&\quad + \langle [u, y]_m, y \rangle \left[\frac{8s y_i y_j}{\alpha^5} - \frac{2}{\alpha^4} (b_j y_i + b_i y_j) - \frac{2s}{\alpha^3} \delta_i^j \right]. \quad (4.14)
\end{aligned}$$

Similarly, by direct computation the value of second term of the equation (4.10) becomes,

$$\frac{\partial^2}{\partial y^i \partial y^j} \left(\frac{\langle [u, y]_m, u \rangle}{2} \right) = 0. \quad (4.15)$$

Hence by substituting equations (4.14), (4.15) into (4.10) we get,

$$E_{ij} = \frac{(n+1)}{2b^2} \left\{ \frac{s}{\alpha(y)} [\langle [u, u_i]_m, u_j \rangle + \langle [u, u_j]_m, u_i \rangle] + \langle [u, y]_m, y \rangle A \right\} \quad (4.16)$$

where

$$A = \frac{8s y_i y_j}{\alpha^5} - \frac{2}{\alpha^4} (b_j y_i + b_i y_j) - \frac{2s}{\alpha^3} \delta_i^j. \quad (4.17)$$

This is the required formula of mean Berwald curvature of homogeneous Finsler-Kropina space. This completes the proof.

5. CONCLUSION

Kropina metrics seem to be the simplest nontrivial Finsler metrics with many interesting application in physics, electron optics with a magnetic field, dissipative mechanics and irreversible thermodynamics [14]. Also, they have interesting applications in relativistic field theory, control theory, evolution and developmental biology. In this article we consider the homogeneous Finsler space with kropina metric. It is an important problem to compute the geometric quantities such as curvatures of homogeneous spaces.

S-curvature is an important geometric quantity in Finsler geometry in that it has some mysterious interrelations with other quantities such as flag curvature, Ricci scalar, etc. Shen showed that the Bishop-Gromov volume comparison theorem holds for Finsler spaces with vanishing S-curvature. Therefore, it is also significant to characterize homogeneous Finsler spaces with vanishing S-curvature. In this paper, we derived an explicit formula of the S-curvature of homogeneous Finsler space with Kropina metric and proved that a homogeneous Kropina space with almost isotropic S-curvature must have vanishing S-curvature. We also derived an explicit formula of the mean Berwald curvature E_{ij} of homogeneous Finsler-Kropina space.

REFERENCES

1. P. L. Antonelli, R. S. Ingarden, and M. Matsumoto, *The Theory of Sprays and Finsler Spaces with Applications in Physics and Biology*, FTPH Vol. 58, Kluwer Academic Publishers, (1993).
2. S. Bacso, X. Cheng and Z. Shen., *Curvature Properties of (α, β) -metrics*, Adv. Stud. Pure Math. Soc., Japan (2007).
3. S. S. Chern, *Finsler geometry is just Riemannian geometry without the quadratic restriction*, Notices Am. Math. Soc. **43**(9), (1996) 959-963.
4. S.S. Chern, Z. Shen, *RiemannFinsler Geometry*, World Scientific Publishers, (2004).
5. X. Cheng and Z. Shen, *A class of Finsler metrics with isotropic S-curvature*, Israel J. Math. **169**, (2009), 317-340.
6. X. Cheng and Z. Shen, *On Douglas metrics*, Publ. Math. Debrecen **66**, (2005), 503-512.
7. S. Deng and Z. Hou, *The group of isometries of a Finsler space*, Pac. J. Math. **207**, (2002), 149-157.
8. S. Deng and X. Wang, *The S-curvature of homogeneous (α, β) -metrics*, Balkan Journal of Geometry and Its Applications, Vol.15, No.2, (2010), pp. 39-48.
9. S. Deng and Z. Hou, *Homogeneous Finsler spaces of negative curvature*, Journal of Geometry and Physics, 57, 657-664, (2007).
10. Z. Hu and S. Deng, *Three dimensional homogeneous Finsler manifolds*, Math. Nachr. 285, No. 10, (2012), 1243-1254.
11. S. Kobayashi and K. Nomizu, *Foundations of Differential Geometry*, Interscience Publishers, Vol. 1, 1963; Vol. 2, (1969).
12. V. K. Kropina, *On projective Finsler spaces with a certain special form*, Nauch. Doklady vyss. Skoly, fiz.-mat. Nauki, Russia, 1959(2), (1960), 38-42.
13. Shaoqiang Deng, *The S-curvature of homogeneous Randers spaces*, Differential Geometry and its Applications, **27**,(2009), 75-84.
14. C. Shibata, *On Finsler spaces with Kropina metric*, Rep. Math. Phys., **13**, (1978), 117-128.
15. Z. Shen, *Volume comparison and its applications in RiemannFinsler geometry*, Adv. Math. **128**, (1997), 306-328.

¹ DEPARTMENT OF MATHEMATICS, KUVEMPU UNIVERSITY, JNANA SAHYADRI, SHANKARAGHATTA-577 451, KARNATAKA, INDIA.

E-mail address: kchandru2510@gmail.com

² DEPARTMENT OF MATHEMATICS, KUVEMPU UNIVERSITY, JNANA SAHYADRI, SHANKARAGHATTA-577 451, KARNATAKA, INDIA.

E-mail address: nmurthysk@hotmail.com